Krzysztof Wiśniewski

Kaon and Antikaon production in Heavy-Ion Collisions for the reactions Ru+Ru and Ru+Zr at 1.69 AGeV beam kinetic energy

PhD Thesis supervised by
dr hab. Krystyna Siwek-Wilczyńska

Institute of Experimental Physics
Warsaw University

June 2000

Contents

1 Introduction **1**

2 Experiment **7**

2.1 Beam and target 7

2.2 The FOPI detector setup 8

2.2.1 Phase 1 9

2.2.2 Phase 2 10

2.2.3 Helitron 14

2.2.4 Geometrical acceptance 15

3 Event characteristics **17**

3.1 Event Selection 17

3.2 Event centralities 21

3.3 Reconstruction of the reaction plane 23

4 Distributions of pions, protons and deuterons **27**

4.1 Identification 27

4.2 Momentum-space distribution 28

4.3 Kinetic-energy spectra 28

4.4 Directed side-flow pattern of protons 30

5 Identification of K^+ mesons **33**

5.1 CDC-Barrel matching 34

5.2 Charge and mass determination 35

5.3 Identification criteria 36

5.4 Efficiency of the K^+ meson identification procedure 39

5.5 Background content in the K^+ meson yields 43

5.6 Systematic uncertainty of K^+ meson yields 45

6 K^+ meson phase-space distribution **47**

6.1 Acceptance 47

6.2 Transverse-mass spectra 48
6.3 Apparent temperatures 52
6.4 Rapidity-density distribution 52
6.5 Centrality dependence of the K^+ mesons production probability . . . 54
6.6 Thermal equilibrium 57
6.7 Kinetic-energy spectra 58
6.8 Chemical equilibrium 60
6.9 Comparison to predictions of the transport model RBUU 60

7 Directed side-flow pattern of K^+ mesons 63
7.1 Flow pattern 63
7.2 Centrality dependence of the K^+ meson flow 64
7.3 Comparison to predictions of the RBUU transport model 65
7.4 Dependence of the K^+ meson flow on transverse momentum 66

8 K^- meson phase-space distribution 69
8.1 K^- meson identification 69
8.2 K^- to K^+ meson yield ratio 73
8.3 Comparison to predictions of the RBUU transport model 75
8.4 Influence of the final-state interactions 77

9 Summary 83

A Appendix 85
A.1 Bethe-Bloch formula 85
A.2 Kinematical variables 86
A.3 Radially-expanding, thermally-equilibrated source 87
A.4 RBUU transport model 87

List of Figures

2.1 The FOPI detector setup . . . 9
2.2 Central Au+Au collision . . . 11
2.3 CDC sector . . . 12
2.4 Particle identification with the CDC . . . 13
2.5 Geometrical acceptance of FOPI . . . 15

3.1 Event-selection conditions . . . 18
3.2 PMUL distribution . . . 19
3.3 Rejection of Ru+O reactions . . . 20
3.4 TMUL distribution . . . 21
3.5 Centrality parameters . . . 23
3.6 Reaction plane . . . 25
3.7 Accuracy of the reaction plane reconstruction . . . 26

4.1 Momentum-space distribution of π^- mesons, protons and deuterons . 28
4.2 Kinetic energy spectra of π^- mesons, protons and deuterons . . . 29
4.3 Temperature and radial flow velocity . . . 30
4.4 Directed side-flow pattern . . . 31

5.1 Upper momentum limit for the K^+ meson identification with the CDC 34
5.2 Charge determination with CDC-Barrel combined measurements . . . 35
5.3 Particle identification with TOF measurements . . . 36
5.4 K^+ meson identification with CDC-Barrel combined measurements . . 37
5.5 Track quality . . . 38
5.6 Upper momentum limit for the K^+ meson identification with the CDC-Barrel combined measurements . . . 39
5.7 Efficiency of the particle identification procedure . . . 41
5.8 Efficiency of K^+ meson identification procedure (m_t versus $y^{(0)}$) . . . 42
5.9 Efficiency of K^+ meson identification procedure (E_{cm}^{kin} versus Θ_{cm}) . . 42
5.10 Background content in K^+ meson yields . . . 44
5.11 Systematic errors . . . 46

6.1 Momentum-space distribution of K^+ mesons 48
6.2 K^+ meson yields as a function of transverse mass 49
6.3 Systematic errors of K^+ meson yields 50
6.4 Transverse-mass spectra of K^+ mesons 51
6.5 Apparent temperatures 52
6.6 Rapidity density distribution of K^+ mesons 53
6.7 Centrality dependence of K^+ meson $dN/dy^{(0)}$ distributions 55
6.8 Centrality dependence of K^+ meson production probability 56
6.9 Centrality dependence of kaon production in Ni+Ni collisions at 1.8 A GeV 57
6.10 K^+ meson production in Au+Au collisions at 1.0 A GeV 57
6.11 Production probabilities of strange and non-strange mesons 58
6.12 Kinetic-energy spectra of K^+ mesons 59
6.13 Predictions of the RBUU model 61
6.14 Geometrical bias on rapidity-density distributions 62

7.1 Directed side-flow pattern of K^+ mesons 64
7.2 Centrality dependence of K^+ meson flow 65
7.3 Transverse-momentum dependence of the flow signal 67

8.1 K^- meson identification conditions 70
8.2 Upper momentum limit for the K^- meson identification 71
8.3 K^- meson yield at different momenta 72
8.4 Momentum-space distribution of K^- mesons 73
8.5 K^- to K^+ meson yield ratio 75
8.6 Geometrical bias on particle yield ratio 78
8.7 Influence of the Coulomb force on charged particle spectra (1) 81
8.8 Influence of the Coulomb force on charged particle spectra (2) 82
8.9 Effect of the Coulomb field on K^- to K^+ meson yield ratio 82

A.1 In-medium self energies and effective masses of K^+ and K^- mesons . 90

Chapter 1

Introduction

Heavy-ion collisions at beam kinetic energies between 1 and 2 GeV per atomic mass unit are very complex processes. It is commonly believed that in the early phase of a central collision a highly dense and hot system is formed. The matter is compressed to a density which exceeds 3 times the normal nuclear matter density ($\rho_0 \simeq 0.16 \frac{\mathrm{nucl.}}{\mathrm{fm}^3}$), and is heated to a temperature in the order of 100 MeV [1]. About 10 fm/c ($3.3 \cdot 10^{-23}$ s) after the beginning of the reaction, the compression energy starts being released in the form of a collective expansion of the system. This leads to a so-called freeze-out of the participating matter after about 30 fm/c. The available energy is sufficiently high so that internal degrees of freedom of nucleons are excited, and other baryons and mesons are produced.

It is very important that the system which is formed during the collision stays in the hot and dense state for a considerably long period of time, longer than the strong-interaction time scale $\simeq 10^{-24}$ s. Thus, for example, heavy-ion experiments are uniquely suitable for studying the compressibility of the nuclear matter far from its ground state (ρ_0, T=0) and for testing parameters of the nuclear equation of state (EOS).

Yet, due to their complexity, heavy-ion experiments are difficult to carry out and interpret. Often conclusions can be drawn only from comparing the experimental results to the predictions of the microscopic, non-equilibrium transport models, like Quantum Molecular Dynamics (QMD) [2, 3] or Boltzmann-Uehling-Uhlenbeck (BUU) [4, 5] models.

Another major problem is the lack of observables that reflect directly the collision scenario, especially its early phase, when the high density is reached, and the most interesting phenomena take place. The weakly interacting probes (leptons) can escape the dense zone undisturbed, but their abundances are scarce. In spite of that, a few experiments aim (or will aim) at the detection of the weakly interacting products of the heavy-ion reactions [6, 7, 8].

The hadronic matter, produced very abundantly during the collisions, interacts strongly. Thus, the information possibly carried out by the hadronic probes is obscured by the copious final-state interactions. Nonetheless, recently there has been a considerable effort, both experimentally and theoretically, to investigate the production of *strange* hadrons, among which K^+ and K^- mesons were of particular interest.

The K^+ meson consists of the $u\bar{s}$ pair, whereas the K^- meson is made of $\bar{u}s$ pair. The threshold kinetic energy for the production of K^+ mesons in an elementary, nucleon-nucleon collision on a stationary target, $NN \rightarrow K^+YN$, is of the order of 1.6 GeV (Y being a strange hyperon Λ or Σ). The threshold kinetic energy for the K^- mesons production in the process $NN \rightarrow NNK^+K^-$ is around 2.5 GeV.

Many predictions based on transport-model calculations show consistently that at beam energies in the range from 1 to 2 A GeV, which are near or below the threshold energies, strange mesons are produced in the early phase of the collisions, in the dense and hot reaction zone. The dominant production channels go via many-step processes, and involve intermediate baryon resonances and/or pions [9]. This interesting hypothesis is supported by the experimental observation that there are more K^+ mesons produced per participant in central than in peripheral Au+Au collisions at 1 A GeV beam kinetic energy [10].

Strange particles are in a sense different from other hadronic probes. They contain the s quark. Strangeness is conserved in the strong interactions. Thus, for example, strange and antistrange particles are produced together, like in the elementary processes mentioned above. Strange particles decay weakly. The mean life time of both K^+ and K^- mesons is 12.37 ns, which corresponds to $c\tau$ of about 3.7 m. The most important decay modes are the leptonic one into the (μ, ν_μ) pair (63.5% branching ratio) and the hadronic one into the (π, π^0) pair (21.2% branching ratio) [11].

Once produced, the strangeness can rarely be re-absorbed during the collision. For a strong interaction this would require a collision of particles that contain the s and $\bar{s}$ quarks, which is not very likely. On the other hand, weak processes, which may not conserve strangeness, occur very rarely by themselves.

There is, however, a tremendous difference between the interactions of K^+ and K^- mesons in the hadronic matter, especially at low relative momenta [12]. K^- mesons can easily interact with nucleons to form Λ or Σ resonances, of which the strangeness content is the same as that of the K^- mesons (S=-1). Therefore, the cross section for the reaction $K^-N \rightarrow \pi Y$ is relatively high – in vacuum, $\sigma \simeq 45$–100 mb for momenta lower than 0.7 GeV/c [13]. Due to the anti-strange quark content, K^+ meson interact mainly elasticly. For momenta lower than 0.7 GeV/c the cross section of the elastic scattering $K^+N \rightarrow K^+N$ is in the order of $\simeq$10 mb [14]. This corresponds to a relatively long mean-free path, which at the normal nuclear matter density would be about 5-7 fm. The dominant charge-exchange process $K^+n \rightarrow K^0p$ has a 2–10 times smaller cross section [15]. Therefore, particularly K^+ mesons are considered to be penetrating probes and a promising tool to infer information about the conditions reached in the early phase of a collision.

First heavy-ion experiments that aimed at studies of strange-particle production slightly above the threshold energies were performed in the early 80's in Berkeley and in Dubna. The Λ hyperon production was investigated with streamer chambers [16, 17, 18], whereas the production of K^+ mesons was studied with magnetic spectrometers [19, 20]. Experiments on subthreshold production of K^- mesons were performed at BEVALAC [20] and, later, with the Fragment Separator at SIS [21]. Already at that time, an enhanced production rate of K^- meson was observed as compared to theoretical predictions.

In heavy ion collisions, the abundances of produced particles depend directly on the bulk properties of the nuclear matter, the EOS, and the collision scenario. Already in early theoretical works [22], it was suggested that, depending on the parameters of the EOS, the K^+ meson yield in central Au+Au collisions at 0.7 A GeV beam kinetic energy can change 2-3 times. A systematic comparison of the K^+ meson production rates for systems of different sizes and energies to the predictions of the QMD transport model favoured the soft EOS, with a small value of the incompressibility parameter, $\kappa \simeq 200$ MeV [23, 24, 25]. The discussion presented recently in [26] addressed the question of thermal equilibrium of the reaction products. It was argued that the final-state momentum-space distribution of K^+ mesons in Ni+Ni collisions at 1.93 A GeV beam kinetic energy is compatible with an isotropic emission pattern with a radial flow component. In contrast, the results obtained for reactions at higher beam energies suggested a sizable anisotropy due to the longitudinal expansion of the system [27].

In this work greater emphasis is put on another subject connected to the production of strange mesons, for which heavy ion experiments offer excellent testing grounds. The studies of in-medium modifications of hadron properties, their effective masses and interactions, were triggered by the idea of the breakdown and the restoration of the chiral symmetry of Quantum Chromodynamics (QCD) [28, 29, 30]. In the low energy limit, the properties of QCD are defined by the symmetries of its Lagrangian. In a world of an exact chiral symmetry, all the quarks should be massless. From the hadron masses, which are evidently not equal to zero, it is obvious that this state is not realized by nature and that the QCD Lagrangian contains terms which are not chirally symmetric. These terms, with the light quark masses in the order of : $m_u \simeq 5$ MeV/c^2, $m_d \simeq 9$ MeV/c^2, $m_s \simeq 175$ MeV/c^2 [31, 32], mix the left- and right-handed quarks, and break the symmetry of QCD explicitly. Moreover, since the hadron-mass spectrum does not exhibit systematic degeneracy with respect to parity, it is believed that the symmetry of QCD is broken spontaneously as well. However, it is also anticipated that the spontaneously broken symmetry of QCD is partially restored in a compressed and hot baryonic matter [29, 33]. And, as the symmetry governs the properties of hadrons, its restoration should manifest itself in modifications of hadron interactions and effective masses in a dense and hot environment [34, 35], such as that reached in heavy-ion collisions.

The in-medium modifications of kaons were first investigated theoretically in [36, 37]. Since then, it has become questionable whether these effects originate directly from the restoration of the chiral symmetry, as anticipated hitherto [34]. Nonetheless, at present, a wide variety of theoretical approaches still predict consistently that the K^+-nucleon interaction should become slightly more repulsive and the K^--

nucleon potential strongly attractive in the dense nuclear environment [39]. In other words, it is expected that in the medium the kaon effective masses change. The K^+ meson mass increases slightly, while the K^- meson mass drops considerably with increasing baryon density. This phenomenon is of particular interest, as it would lead to the formation of a kaon condensate in a dense nuclear matter [36, 37], which, in turn, would have consequences for the evolution of neutron stars. As pointed out in [40], stars with masses smaller then previously estimated would collapse into black holes.

The evidence for the modifications of kaon properties at low densities is based on the analysis of K^+-nucleus scattering [41] and on the studies on kaonic atoms [42]. The results of K^+-nucleus scattering experiments favour moderate repulsion of the K^+N interaction at the normal nuclear matter density (ρ_0), which is what one expects on grounds of K^+-nucleon scattering lengths [43, 44]. On the other hand, the results on kaonic atoms imply considerable attraction of the K^-N interaction at ρ_0, whereas from the empirical K^--nucleon scattering lengths [43] it is expected that the K^-N interaction is repulsive. As pointed out in [38], this transition, from repulsion in vacuum to attraction at non-vanishing densities, can be understood as an effect of the in-medium modifications of the $\Lambda(1405)$ properties.

In heavy-ion collisions higher densities are reached. Therefore, the in-medium effects can be probed in conditions in which they are thought to be much more pronounced. One expects that their influence on kaons is twofold.

First of all, because the kaon effective masses are modified, the production threshold energies change. It means a small reduction of the production rate of K^+ mesons, of which the effective mass is supposed to grow slightly, and a substantial rise of the production rate of K^- mesons, the effective mass of which is thought to drop considerably. This hypothesis was confirmed experimentally by the results of the KaoS collaboration [45]. The K^+ and K^- meson yields were measured at around mid-rapidity in Ni+Ni collisions at two beam energies equally distant from the threshold energies for the production of mesons in elementary, NN collisions. The production cross sections of both meson species turned out to be similar at these equivalent energies, whereas the respective production rates in NN collisions slightly above the production thresholds differ by one order of magnitude. This intriguing result, which indeed suggested the changes of the kaon effective masses and production threshold energies, needs to be confirmed by independent measurements, possibly in different regions of the momentum space. However, it was suggested recently that in heavy ion experiments, the chaotic as well as the coherent movement of the baryonic matter may additionally mask the effect of the in-medium potentials on the measured antikaon yields [46].

Secondly, the effective kaon-nucleon potentials change as kaons propagate in the nuclear medium of varying density. The gradients of potentials, i.e., forces acting on kaons, are thought to modify their final-state phase-space distributions. In particular, K^+ mesons are expected to be pushed away from the regions of high baryon density by a slightly repulsive potential, while K^- mesons are thought to be attracted towards the dense collision zone by a strongly attractive force. In order to investigate this problem, the FOPI collaboration measured the directed side-flow pattern of K^+ mesons in the Ni+Ni system at 1.93 A GeV beam kinetic energy [47]. It was shown that K^+ mesons do not flow in the direction of baryons. This is surprising because the flow of Λ hyperons, which are produced together with K^+ mesons, clearly follows the proton flow. The observation was interpreted as a hint for additional repulsion felt by K^+ mesons during the propagation through the medium. Comparison of the result to the predictions of transport models revealed that it can be understood as an effect of the in-medium modifications of the K^+N potential [48]. However, it was pointed out that if the momentum dependence of the K^+N interaction is properly taken into account, the sensitivity of the observable to in-medium effects gets considerably reduced [49]. It was also suggested that the effect may depend strongly on the lifetimes of the nuclear resonances which are used in the model calculations [50]. Recently, the azimuthal anisotropy of the K^+ meson phase-space distribution was observed in the Au+Au system at 1 A GeV beam kinetic energy [51]. The results were compared to the predictions of a transport model, and again interpreted as a hint for the in-medium modifications of kaons. Certainly more experimental evidence is needed in order to elucidate the outstanding questions.

In this work, the measurements on K^+ and K^- mesons in $^{96}Ru+^{96}Ru$ and $^{96}Ru+^{96}ZrO_2$ collisions at 1.69 A GeV beam kinetic energy are reported. The data were taken in 1996 with the FOPI detector system, installed on the beam of SIS at GSI/Darmstadt. The question of the in-medium modifications of kaons is addressed by comparing the experimental results to the predictions of the RBUU transport model of the Giessen group. Some detailed information about the model is given in Appendix A.4.

The work is organized in the following way. In Chapters 2 and 3, the details about the experiment, about the detector setup, and about the data analysis are given. In Chapter 4, the method and results of extracting the freeze-out conditions by analysis of pion, proton, and deuteron spectra is presented. In Chapter 5, the kaon identification procedure is described. In Chapter 6, the final-state momentum-space distribution of K^+ mesons and the centrality dependence of the K^+ mesons production are discussed. It is shown that the accuracy of the measurements does not allow to draw firm conclusions about the in medium effects on kaons. On the other hand, the results are exploited for studying the thermal and chemical equilibrium of the system. In Chapter 7, the directed side-flow pattern of K^+ mesons and the dependence of the measured signal on the transverse momentum are discussed. The comparison of the results to the predictions of the RBUU transport model reveals that there is a potentially high sensitivity of these observables to the in-medium effects on kaons. The experimental observations are best understood if the modifications of the K^+ meson effective mass are taken into account in the model. In Chapter 8, the K^- to K^+ meson yield ratio is studied. It is shown that the yield ratio varies across the measured fraction of the momentum space. It is argued that the observation can neither be explained as the effect of the final-state interactions nor can it have a purely kinematical origin. The experimental results can be reproduced by the RBUU transport model only if the in-medium modifications of kaons are considered. In Chapter 9, the concluding remarks are given.

Chapter 2

Experiment

2.1 Beam and target

The accelerator facility at Gesellschaft für Schwerionenforschung (GSI) in Darmstadt, which includes, among others, the Schwerionensynchrotron (SIS), nowadays delivers beams of light and heavy ions as well as protons and pions. In the case of ion beams, the available beam kinetic energy per atomic mass unit is described by the formula

$$E_{\mathbf{beam}}/A = m_{\mathbf{proton}}c^2 \left(\sqrt{1 + \left(\frac{B\rho \cdot c}{m_{\mathbf{proton}}c^2} \frac{Q}{A} \right)^2} - 1 \right) . \tag{2.1}$$

Q and A denote the charge and the mass number of beam ions. $B\rho$ is the rigidity of the synchrotron. The maximum value for the rigidity of SIS equals 18.7 Tm. Hence, the available beam kinetic energies are as high as 2 GeV per nucleon.

In September 1996, a series of experiments was performed at SIS with the beams of $^{96}_{44}Ru$ and $^{96}_{40}Zr$ isotopes on $^{96}_{44}Ru$ and $^{96}_{40}Zr^{16}_{8}O_2$ targets. There were two primary goals of these experiments :

- investigation of transparency of the nuclear matter and its dependence on the collision energy,
- investigation of strange meson production.

According to formula 2.1, the available beam kinetic energy grows with the charge of the beam ions. On the other hand, the beam intensities decrease with increasing ion charge. For the Ru beam, the highest charge state that could be used in order to keep a sufficient beam intensity was the 42+ charge state. The resulting beam kinetic energy was 1.69 A GeV. The beam energy was slightly lower (1.58 A GeV) in the case of the Zr beam because the highest used charged state was 41+.

At about 2 A GeV beam kinetic energy, the production rate of K^+ mesons is in the order of 1 per 10 central collisions. It decreases rapidly with decreasing beam energy. The dependence of the production cross section of K^+ mesons on the beam kinetic energy can be approximated by the formula [26]

$$\sigma(\mathrm{K}^+) \simeq (E_{\mathrm{beam}})^{5.3} \; . \tag{2.2}$$

At the same energy, the production cross section of K^- mesons is typically an order of magnitude lower [45]. Therefore, in this work, which concentrates on the production of both K^+ and K^- mesons, only reactions with the $^{96}_{44}$Ru beam at the highest kinetic energy of 1.69 A GeV were investigated. The production rate of K^- mesons was too low in the case of the reactions at lower beam energies.

Two target foils were used :

- 430 mg/cm^2 of Ru,
- 380 mg/cm^2 of ZrO_2.

Both target thicknesses were chosen to assure about 1% probability of reactions of the beam ions with the Ru and Zr nuclei from the target. The beam spills were typically 10–12 second long. In order to keep the reaction rate during the spills below $5{\cdot}10^2$ Hz (as required by the operation conditions of the detector apparatus), the beam intensity was kept below $5{\cdot}10^4$ ions per second. This corresponded to an average time interval between two incoming ions in the order of 20 μs. In these conditions, the average time between two reactions was about 2 ms.

The target foils were placed inside the FOPI detector setup, which will be described in greater detail in the following section.

2.2 The FOPI detector setup

The FOPI detector system [52, 53] is a general purpose, modular setup of several subdetectors, used for fixed target experiments. It has a nearly complete azimuthal symmetry and almost full geometrical acceptance[1]. A schematic view of the detector is shown in Figure 2.1.

The FOPI system is capable to detect simultaneously charged reaction products: heavy fragments of Z up to 20, light fragments (protons, deuterons, tritons) and mesons (pions, kaons, etc.). Particles are identified and their four-momenta are determined by the measurements of the specific energy loss in the detector material (dE/dx), the track curvature radius in the magnetic field (ρ), and the velocity (v), derived from the information on time-of-flight from the target position to the detector (TOF). In addition, shortly-lived neutral particles are identified by invariant mass reconstruction of their decay products.

[1] The name "FOPI" is an abbreviation of "*FOur PI*", and originates from the large, nearly completely 4π geometrical acceptance of the setup.



Figure 2.1: *The FOPI detector setup.*

2.2.1 Phase 1

The forward part of the FOPI setup, i.e., the so-called Phase 1, was completed in 1990 [52]. It consists of two systems of scintillating strips: the Zero Degree (ZD) and the Plastic Wall (PLAWA) detectors. The ZD detector allows the detection of charged particles emitted from the target position at laboratory polar angles (Θ_{lab}) between 1.2° and 7.5° over the full azimuthal range. The PLAWA detector covers the full azimuth of the polar angles from 7° to 30°. Both detectors provide the information about the TOF and the energy deposited by particles in the scintillator material (ΔE).

Identification of particles is possible for sufficiently high velocities, higher than the punch-through velocity. In such cases, the information about ΔE allows to determine the specific energy loss, dE/dx, of a particle. In order to provide the information on dE/dx for particles of which velocities are lower than the punch-through velocity, the Parabola (ionization chamber) and the Rosace (thin scintillator wall) were usually placed in front of the PLAWA and the ZD detectors respectively[2].

The information on dE/dx and v allows to derive easily particle charge but not particle mass (cf. formula A.1, and notice that the dE/dx of a particle passing through matter does not depend explicitly on its mass). Therefore, Phase 1 detectors are capable (only) of particle identification with respect to their charge. A series

[2]Since 1994, when the forward drift chamber, Helitron, was introduced to the detector setup, Parabola has no been used any more.

of successful experiments which focused on the production of nuclear fragments followed the completion of the Phase 1 system [54, 55, 56, 57, 58]. For almost all practical purposes of those experiments, an estimate that the mass of a fragment is given by $M_{fr}= 2\times Z_{fr}$ was sufficiently accurate.

Unfortunately, PLAWA and ZD detectors can not be used for the identification of K^+ and K^- mesons, which requires particle mass determination. Therefore, in this work, the information from Phase 1 detectors is used only for determining global event characteristics, like collision centrality or orientation of the reaction plane (see Sections 3.2 and 3.3 for details).

2.2.2 Phase 2

Phase 2 of the FOPI setup [53] was designed for experiments with beams of higher kinetic energies, up to maximal energies delivered by the SIS. Only lighter nuclear fragments survive such high-energy collisions. Moreover, the internal degrees of freedom of nucleons participating in the reactions are excited, and particles like pions, kaons and antikaons are produced. For their identification the information about their mass, not only charge, is essential. Phase 2 of FOPI was built to meet the requirement of particle mass determination. Its completion was followed by experiments which focused mainly on the production of mesons [47, 59, 26, 60].

Phase 2 of FOPI consists of the tracking device – Central Drift Chamber (CDC) – and the Plastic Scintillator Barrel (cf. Figure 2.1). Both detectors operate in the magnetic field of 0.6 T, homogenous and parallel to the beam axis (z), which is produced by the superconducting solenoid. The Barrel surrounds the CDC and provides additional reliability of the particle identification. Detailed information about the operation of the CDC and the Barrel as well as about the calibration procedures can be found in PhD theses [61, 62]. Here only the most important technical information will be given.

Up to 100 particles per central heavy-ion collision can make their way through the CDC. Figure 2.2 shows how such an event looks like in the plane perpendicular to the beam axis (xy). Dots represent individual hits from which particle tracks are reconstructed. The amount of energy lost by particles due to ionization is illustrated by the sizes of the dots. Trajectories of negatively and positively charged particles, depicted by the lines, are bent in opposite directions in the magnetic field. An average hit multiplicity per particle trajectory is about 50. Therefore, as many as 5000 hits can be registered in a single event. A correct assignment of hits to an appropriate trajectory is a serious challenge for the tracking algorithm, of which several different types are used [61, 63, 64]. Matching between tracks reconstructed in the CDC and hits registered in the Barrel [61, 62] (filled squares in Figure 2.2) brings in additional difficulty, but is necessary for identification of rare particles.



Figure 2.2: *Trajectories of charged products of a central Au+Au collision at 1 A GeV beam kinetic energy, as seen by the CDC and Barrel in the plane perpendicular to the beam axis. About 80 particle trajectories are reconstructed out of nearly 3800 hits.*

CDC

The CDC covers the full azimuth of the polar angles between 30° and 150°. It allows the tracking of charged particles at radial distances between 20 and 80 cm from target position. It is subdivided into 16 sectors in the xy plane. The view of one of the sectors in the xy plane is shown in Figure 2.3. In the middle of the sector, an anode plane is formed by 60 grounded read-out wires. A homogenous electric field of 750 V/cm is provided by the so-called drift wires located on the border of the sector and on the inner and outer radii of the chamber. In between the read-out wires, the potential wires, on -1.5 kV voltage, are located. They provide a strong electric field in the vicinity of the read-out wires so that charge multiplication can take place.

While passing through the CDC, charged particles ionize the gas, typically a mixture of 88% Argon, 10% isobutane, and 2% methane under the atmospheric pressure. The produced electrons drift towards the anode planes on straight trajectories, determined by the interplay between the electric and magnetic fields, with velocity in the order of 45 μm/ns. The maximum drift time of the electrons is about 5 μs. In the vicinity of each read-out wire, before the charge is collected, the multiplication of the charge occurs ($\times 10^3$). The signals are read on both ends of the read-out wires with a system of ADC's, which allows to digitize the complete signal shape over a 10 μs time interval.

The hit coordinates, i.e., the location at which the primary ionization took place, can be calculated back. From the arrival time of the signals on both ends of the wire, the hit position in the xy plane is derived with the accuracy of 300 μm. From the charge division between both ends of the wire, the hit coordinate along the z



Figure 2.3: *A schematic view of one sector of the CDC in the plane perpendicular to the beam axis. The insert gives the magnified view of the anode plane in the middle of the sector. Staggering of the read-out wires on both sides of the anode plane can be observed.*

axis is deduced with a 5 cm accuracy. The energy loss of particle can be derived from the total charge collected on the wire, which is proportional to the primary ionization.

The tracking program groups individual hits to form particle trajectories, which are fitted with a helix curve. For each trajectory the curvature radius (ρ), the direction of emission (Θ_{lab} and ϕ_{lab}), and the mean value of the energy loss (dE/dx) are determined. The accuracy of dE/dx determination is momentum dependent. It varies form 5 to 15% with momenta increasing from 0.1 to 2 GeV/c.

The ratio of transverse momentum over charge of a particle is derived via formula

$$\frac{p_t}{Q} = B\rho \ . \tag{2.3}$$

The accuracy of p_t determination in the order of 5% to 15% is achieved.

Particle identification with the CDC

Figure 2.4a shows the identification plot in the form of a correlation matrix between the logarithm of dE/dx and the ratio p/Q of particles for which trajectories were reconstructed within the CDC. The data comes from the Ru+Ru experiment at 1.69 A GeV beam kinetic energy. The lines plotted on top of the data show the dependence of the specific energy loss on particle momenta, calculated according to the Bethe–Bloch formula A.1[3]. Figure 2.4b shows the mass spectrum, i.e., the projection of the matrix from panel (a) on the mass parameter in the Bethe–Bloch formula. The mass parameters are multiplied by -1 for particles of which trajectories have opposite curvature than proton trajectories. The plotted quantity is often called $Mass_{CDC}$ since its determination is based on the information delivered by the CDC only. The plot demonstrates a good separation between different particle

[3]Some modifications of formula A.1 are needed in order to account for non linear response of the detector.



Figure 2.4: *Particle identification with the CDC. The data come from the Ru+Ru experiment at 1.69 GeV beam kinetic energy. (a) The matrix of correlations between energy loss dE/dx and ratio p/Q of particles. (b) The mass spectrum obtained by projection of the matrix from panel (a) on the mass parameter in the Bethe–Bloch formula.*

species: pions, protons and deuterons, which, however, gets worse with increasing momentum (cf. Figure 2.4a). Particles can be selected according to the values of the $Mass_{CDC}$ parameter. Studies on pions and on light fragments (p, d, t) are often performed using the information delivered by the CDC only.

Plastic Scintillator Barrel

The CDC is surrounded by the Plastic Scintillator Barrel, a set of 180 individual scintillator strips which form a cylinder of a 111 cm radius (cf. figures 2.1 and 2.2). The strips are 240 cm long, and have a 4×3 cm rectangular shape in cross section. It was originally planed that the polar-angle range covered by the Barrel matches that of the CDC. However, because of the high double-hit rate in the scintillator strips, the Barrel was moved backwards so that the opening angle, as seen from the target position, is currently $40° < \Theta_{lab} < 140°$. The Barrel covers about 85% of the full range in the azimuthal angle. Only two sets, of six strips each, are missing due to technical reasons (cf. Figure 2.2).

Scintillation light produced in a given strip by charged particles is collected on both strip ends by photomultipliers[4]. Both the light output and the arrival time of the signal are registered. The time is measured against the signal from the start detector (see Section 3.1 for more information). The delay due to the passage of the beam ions from the start detector to the target is subtracted.

[4]The role of the light guides which connect the scintillator strips with the photomultipliers is not discussed in this work.

The TOF of a particle from the target position to the point of the detection is derived via the formula

$$TOF = \frac{T_L + T_R}{2} - \frac{L}{2 \cdot v_{light}} , \tag{2.4}$$

where T_L and T_R denote the arrival time of the signal at the upstream and at the downstream ends of the strip respectively, and L=240 cm is the strip length. The accuracy of the TOF determination varies from strip to strip but is in the order of 300–400 ps.

The distance from the middle of the strip to the position at which a particle crossed the strip along the z axis (hit position) is derived via the formula

$$Z = \frac{T_L - T_R}{2} v_{light} . \tag{2.5}$$

$v_{light} \simeq 16$ cm/ns denotes the velocity of the light-pulse propagation in the scintillator material. The Z coordinate of the hit is determined with 8 cm accuracy, which is limited by the accuracy of time measurements. The hit position in the xy plane is given by the azimuthal angle ϕ of the strip with accuracy of 2°, limited by the dimensions of the strip.

The velocity of the particle is derived from the position of the hit and the TOF information via the relation

$$v = \frac{s}{TOF} = \frac{(2 \cdot \rho \cdot \arcsin(\frac{R_B}{2 \cdot \rho}))/\sin{(\Theta_{lab})}}{TOF} , \tag{2.6}$$

where s is particle flight path, ρ is curvature radius of particle trajectory in the magnetic field, Θ_{lab} is the emission polar angle, and R_B=111 cm is the radius of the Barrel.

In case if two or more adjacent strips of the Barrel register hits, the correspondence between TOF and the Z measurements on the subsequent strips is checked. When the differences in TOF and in the hit position along the z axis (ΔTOF and ΔZ) are smaller than 300 ps and 25 cm respectively, it is assumed that the hits originate from one particle, of which the trajectory is sufficiently bent so that it passed through several Barrel strips. The hit position and the TOF for this particle is taken from the strip which registered the highest energy deposited by the particle.

The operation of the CDC and the Barrel is essential for the analysis on which this work focuses. The procedure of particle identification with CDC–Barrel combined measurement will be described in greater detail in Chapter 5.

2.2.3 Helitron

The FOPI Phase 2 is completed by the Helitron (cf. Figure 2.1). It is the forward drift chamber, which will eventually play the same role as the CDC does in the backward hemisphere. It will add information about the mass to the charge derived



Figure 2.5: *The geometrical acceptance of the FOPI detector setup for a mass symmetric collision of a beam at 1.69 A GeV kinetic energy with a fixed target located in the middle of the setup. (a) The solid lines are drawn for kaons of momenta* p_{lab}*=0.2, 0.4, 0.6, 0.8 and 1.0 GeV/c. (b) The solid lines show different emission angles from* Θ_{cm}*=20° to* Θ_{cm}*=160° in 20° intervals. The dashed lines correspond to kaon kinetic energies in the center-of-the-momentum reference frame of* E_{cm}^{kin}*=0.2, 0.4, 0.6, 0.8 and 1.0 GeV.*

from the measurement by the Plastic Wall for particles emitted in the mid-rapidity region and at low transverse momenta. Most of the particle yields populate exactly this region of the momentum space. Therefore, it will be of high importance to combine the yields measured by the CDC in the backward direction with those estimated from the Helitron data. Among others, the effort to identify K^+ and K^- mesons with the Helitron is currently being pursued [65] but it is not described in this work. Here, no use is made out of the information from the Helitron.

2.2.4 Geometrical acceptance

In Figure 2.5a the geometrical acceptance of the individual subcomponents of the FOPI detector setup is shown for a mass-symmetric reaction of a beam at a kinetic energy of 1.69 GeV per nucleon. The acceptance is presented in terms of transverse momentum divided by the mass of a particle (p_t/m_0c)[5] versus the rapidity normalized to the rapidity of the center-of-the-momentum (CM) coordinate system $(y^{(0)})$ [6]. Due to the almost full coverage of the available momentum space, in an ideal case, only a small fraction of the charged reaction products will remain not detected by the FOPI setup.

[5]Particle mass is multiplied by the light velocity, c, in order to arrive at a dimensionless quantity.
[6]The definitions of the kinematic variables referred to throughout this work can be found in Appendix A.2.

Chapter 3

Event characteristics

Altogether around 4.8 million events were collected in the Ru+Ru experiment at 1.69 A GeV beam kinetic energy. The number of events registered in the Ru+ZrO_2 experiment at 1.69 A GeV beam kinetic energy was about 3.7 million.

3.1 Event Selection

Rejection of double interactions

From all the subdetectors of the FOPI setup the CDC has the longest operational dead-time. As mentioned in Section 2.2, the ADC system of CDC allows to digitize the complete signal shape over a 10 μs time interval, whereas the maximum drift-time of electrons in the CDC is about 5 μs. When a given event is accepted by the trigger system, it is important to assure that no reactions prior and after this event took place within a 5 μs time period. In an opposite case, a so-called double interaction is encountered and the events are mixed. Since the average time interval between two reactions is in the order of 2 ms, double interactions are in general rare.

In most of the cases, double interactions are rejected already on the trigger level, rejection being based upon the much faster signals from the plastic detectors of the Forward Wall and Barrel. In order to reject double interactions caused by two ions which came within a 100 ns time interval one after the other, the information from the start detector is needed. It is a thin scintillator placed about 2.5 m up-stream from the target. It registers both the arrival time and the energy deposited in the scintillator material by beam ions. Figure 3.1a shows the energy distribution of the signal from the start detector. A double-peak structure can be observed. The peak corresponding to a higher energy deposited by the beam ions (above the minimum in between the two peaks) is attributed to double interactions when two ions pass the start detector very shortly one after the other (within 100 ns interval of the QDC gate). The figure caption gives the condition which was used in order to reject double interaction from subsequent analysis.



Figure 3.1: *(a, b) Time and energy spectra from the start detector. (c) The distribution of the coordinates of the event vertex in the plane perpendicular to the beam axis.*

The start detector delivers also the information about the reference time. Figure 3.1b shows the distribution of the arrival time of the signal from the start detector. The events for which the time was outside a certain limit around the central peak (indicated in the figure caption) were rejected as their reference time was wrong.

Rejection of non-target reactions

The halo detector, which is used to insure the focusing of the beam and as a veto device, consists of two components. The first one is a 2 mm thick scintillator with a cylindrical hole, 7 mm in diameter, placed close to the target. The second part is a 5 mm thick scintillator with a rectangular hole, 10 x 10 mm in cross section, located about 2.5 m upstream from the target, close to the start detector. In normal conditions 98–99% of the ions from the beam pass through the holes of the halo detector.

In order to further suppress the number of reactions that did not occur on target, the coordinates of the event vertex were calculated. First of all, the pairs of tracks in the CDC which intersect were found, and the coordinates of the interception points were calculated. The event-vertex coordinates (x_v, y_v, z_v) were computed as the mean values of the coordinates of the intersection points. The distribution of the event-vertex coordinates in the xy plane with respect to the target position is shown in Figure 3.1c. The caption indicates the conditions which are imposed in order to select only the reactions which took place on target.



Figure 3.2: *The PMUL distributions for minimum-bias (dashed histogram) and medium-bias (dotted histogram) event samples in the Ru+Ru experiment. The sum of the two is depicted by the solid histogram.*

Minimum-bias and medium-bias event classes

Two samples of events were accumulated simultaneously during the experiment. The so-called minimum-bias event sample (MinB) was collected practically without any preselection. The second class, on the other hand, the so-called medium-bias sample (MedB), consisted of events which fulfilled the requirements of some minimum multiplicity of hits registered in the Plastic Wall. The selection was made on the hardware-trigger level. The threshold on the multiplicity was set as high as to select approximately 15% of all the events. Figure 3.2 shows the distributions of multiplicities of particles identified by the Plastic Wall (PMUL) for both considered event samples, collected in the Ru+Ru experiment. Usually some of the hits from the total hit multiplicity in the Plastic Wall can not be assigned to any of the identified particles. Therefore, the PMUL distribution for the MedB event sample shown in Figure 3.2 falls down gradually on the side of low PMUL instead of ending sharply at a certain threshold value.

It was assumed that because of the condition imposed on PMUL, the fraction of central collisions was enhanced in the MedB event sample. Since the production rate of kaons is higher in central than in peripheral collisions, the MedB event sample is of greater interest for the studies which are reported in this work. In order to allow to accumulate more central events within a limited speed of the data acquisition system, the storage rate of MinB events was scaled down by a factor of 32 (cf. Figure 3.2).

The method of selecting events according to their centrality is further discussed in the next section.

Rejection of Ru+O reactions

The rejection of the reactions of the Ru beam ions with the Oxygen nuclei in the ZrO_2 target was based on the multiplicity of detected particles. In Figure 3.3 the



Figure 3.3: *The multiplicity of particles detected by the Plastic Wall (PMUL) versus the multiplicity of particles registered by the CDC (CMUL) in the Ru+Ru (left) and Ru+ZrO_2 (right) experiments. (a,b) The distributions for the minimum-bias event sample. (c,d) The distributions for the medium-bias event sample.*

multiplicity of particles registered by the Plastic Wall (PMUL) is plotted against the multiplicity of particles detected in the backward hemisphere by the CDC (CMUL). In the case of the Ru+Ru experiment (panel a), the more central the collisions are, the higher PMUL and CMUL they have. In contrast, in Figure 3.3b, which shows the multiplicities of particles in the Ru+ZrO_2 experiment, two separate branches can be noticed. The branch characterized by the weaker correlation between PMUL and CMUL corresponds to the reactions of the beam ions with the Oxygen nuclei in the ZrO_2 target. The reactions with Oxygen nuclei can have quite high PMUL, but they are forward peaked. Therefore, their CMUL (which measures the number of particles emitted in backward directions) remains low, even in the case of central collisions. The reactions with Oxygen nuclei in the target can not be, however, easily separated from the more peripheral reactions with Zr nuclei, of which both PMUL and CMUL are low.

Panels (c) and (d) in Figure 3.3 show PMUL plotted against CMUL for the medium-bias event sample in the Ru+Ru and Ru+ZrO_2 experiments, respectively. Because



Figure 3.4: *The distributions of TMUL in the Ru+Ru experiment for minimum-bias (dashed histogram) and medium-bias (solid histogram) event samples. The histogram for the MinB event sample is normalized to the total geometrical cross section of the reaction. The division of the medium-bias sample into three event categories according to collision centrality is depicted with grey shaded areas.*

of the requirement of high PMUL, reactions with Oxygen nuclei can be eliminated from the MedB event sample by requiring CMUL to be higher than 15.

In this work, about 4.38 million and 3.39 million events from the MedB event samples were analysed in the case of the Ru+Ru and Ru+ZrO_2 experiments respectively (cf. Table 3.1). As described, the selection was made according to the conditions imposed on time and energy signals from the start detector, event-vertex coordinates, and multiplicities of detected particles.

3.2 Event centralities

It is commonly accepted that in heavy-ion collisions at intermediate energies the multiplicity of the reaction products is correlated with impact parameter of a collision [66]. In this work, the selection of events according to their centrality is based on the total multiplicity of particles detected by the CDC and by the Plastic Wall,

$$\text{TMUL}=\text{CMUL}+\text{PMUL}\ . \tag{3.1}$$

The dashed histogram in Figure 3.4 shows the TMUL distribution for the MinB event sample weighted with the trigger efficiency. The trigger efficiency was defined by the dead-time of the data acquisition system. It was derived by comparing the number of collected events with the number of collisions expected from the total geometrical reaction cross section (Eq. 3.2). Thus, the dashed histogram in Figure 3.4 is normalized to the total geometrical cross section of the reaction

$$\sigma_{geo} = \pi \cdot (2 \cdot 1.2\ \text{fm} \cdot 96^{1/3})^2 \simeq 3800\ \text{mb}\ , \tag{3.2}$$

where the radius of a ^{96}Ru or ^{96}Zr nucleus is approximated by

$$R_{Ru} = R_{Zr} = 1.2 \cdot 96^{1/3}\ \text{fm} \simeq 5.5\ \text{fm}\ . \tag{3.3}$$

Assuming that there is a monotonic dependence of multiplicity of detected particles on impact parameter [66], the integration of the TMUL distribution from a given value TMUL$_{min1}$ to its upper end gives the value of $\Delta\sigma$ – the centralmost part of the cross section corresponding to the event sample selected with the condition TMUL>TMUL$_{min1}$. Using the geometrical sharp-cut-off model, one can estimate the maximal and the average impact parameters characterizing this event sample

$$b_{geo}^{max} = \sqrt{\Delta\sigma/\pi} \ , \tag{3.4}$$

$$< b >_{geo} = \frac{\int_0^{b_{geo}^{max}} b \cdot 2\pi b db}{\int_0^{b_{geo}^{max}} 2\pi b db} = 2/3 \cdot (b_{max}) \ . \tag{3.5}$$

Using a participant-spectator model, one can subsequently calculate the number of nucleons participating in the reactions. In this model, the participant zone is the overlap volume of two spheres passing each other at a distance b. Each sphere has a radius equal to that of the nucleus $R_{Ru,Zr}$. On these assumptions, the mean number of participants characterizing the sample of events of which impact parameters are less than b_{geo}^{max} is equal to [67]

$$< A_{part} > = \frac{\int_0^{b_{geo}^{max}} (1 + (\frac{3}{\sqrt{2}} - 3)\frac{b}{R_{Ru,Zr}} + (3 - 2\frac{3}{\sqrt{2}})(\frac{b}{R_{Ru,Zr}})^2 + (\frac{3}{\sqrt{2}} - 1)(\frac{b}{R_{Ru,Zr}})^3) \cdot 2\pi b db}{\int_0^{b_{geo}^{max}} 2\pi b db} \ . \tag{3.6}$$

Similarly, one can assign the part of the cross section, $\Delta\sigma$, and the parameters $<b>_{geo}$ and $<A_{part}>$ to the sample of events of which TMUL are in any interval, for example, TMUL$_{min2}$<TMUL<TMUL$_{min1}$. Figure 3.5 shows the values of the parameters determined in this way for events of different TMUL grouped into 10-unit-wide bins.

In this work, the MedB event sample is analysed. The solid histogram in Figure 3.4 shows the TMUL distribution for the MedB event sample weighted with the MedB trigger efficiency. At the upper end, i.e., for very central collisions, where the hardware-trigger selection has no influence, the distribution matches that of the MinB event sample. The grey shaded areas illustrate the division of the MedB sample into three event subsamples which will be referred to in the following part of this work. The events are grouped according to their TMUL values. It is assumed that in this way event categories of different centralities are selected. However, due to fluctuations of TMUL, certainly events of quite different impact parameters may be mixed together in the considered subsamples.

For each subsample, the parameters $\Delta\sigma$, b_{geo}^{max}, $<b>_{geo}$, and $<A_{part}>$ were calculated via the TMUL distribution of the MedB sample shown in Figure 3.4, following the method sketched above. The values of these parameters, the TMUL ranges, and the statistics of the event subsamples are listed in Table 3.1.



Figure 3.5: *(a) The correspondence between the cross section, $\Delta\sigma$, and the limiting TMUL; (b) the dependence of b_{geo}^{max} on TMUL; (c) $<b>_{geo}$ as a function of TMUL; (d) $<A_{part}>$ as a function of TMUL. The values are calculated for 10-unit-wide bins of the TMUL distribution. The dashed lines represent (a) the total geometrical reaction cross section, σ_{geo}=3800 mb, (b) the maximum impact parameter from the geometrical sharp-cut-off model, $b_{geo}^{max} \simeq 11$ fm, and (d) the maximum number of participants, $<A_{part}>=192$.*

3.3 Reconstruction of the reaction plane

The reaction plane is defined as the one that contains the projectile trajectory and the vector of impact parameter $\boldsymbol{b}$ (cf. Figure 3.6a). The method of its determination for each individual event was developed in [68]. Since the reaction products are repelled from the central collision zone in the reaction plane (cf. Figure 3.6), a vector calculated by summing the transverse momenta ($\boldsymbol{p}_t$) of the reaction products in the projectile-rapidity region is supposed to be parallel to $\boldsymbol{b}$. Similarly, a vector calculated from the momenta of the reaction products in the target-rapidity region is antiparallel to $\boldsymbol{b}$. Usually the difference of the two is calculated

$$\boldsymbol{Q} = \sum_n \boldsymbol{p}_t^n \cdot w^n \ . \tag{3.7}$$

TMUL		50 – 85	85 – 95	95 – 140	all
# of events (no selection)	$^{96}Ru+^{96}Ru$	$2.3 \cdot 10^6$	$1.4 \cdot 10^6$	$1.0 \cdot 10^6$	$4.8 \cdot 10^6$
	$^{96}Ru+^{96}ZrO_2$	$1.9 \cdot 10^6$	$1.1 \cdot 10^6$	$0.7 \cdot 10^6$	$3.7 \cdot 10^6$
# of events (after selection)	$^{96}Ru+^{96}Ru$	$1.9 \cdot 10^6$	$1.4 \cdot 10^6$	$1.0 \cdot 10^6$	$4.4 \cdot 10^6$
	$^{96}Ru+^{96}ZrO_2$	$1.6 \cdot 10^6$	$1.1 \cdot 10^6$	$0.7 \cdot 10^6$	$3.4 \cdot 10^6$
$\Delta\sigma$		280 mb	180 mb	120 mb	580 mb
$\Delta\sigma/\sigma_{tot}$		5%	7%	3%	15%
b_{geo}^{max}		4.3 fm	3.1 fm	2.0 fm	4.3 fm
$< b >_{geo}$		3.7 fm	2.5 fm	1.3 fm	2.8 fm
$< A_{part} >$		120	140	170	135

Table 3.1: *The parameters characterizing different event categories selected according to collision centrality given by TMUL.*

The sum runs over the reaction products. The weight w^n can be set to +1 for particles of which rapidities are larger than the mid-rapidity and to -1 for particles emitted in the backward hemisphere.

In this work, the information on the $\boldsymbol{p}_t$ of charged particles is delivered by the Plastic Wall and the CDC. However, as commonly accepted [68], some of the reaction products are excluded from the summation 3.7:

- particles which are emitted within a certain region around the mid-rapidity, $|y^{(0)}|<0.3$, because the direction of their emission has little correlation with the reaction plane,
- pions identified in the CDC, because (as it is also the case for other particles produced during the collision) their correlation with the entrance channel, which defines the reaction plane, is not direct.

For each individual event, the azimuth of the vector $\boldsymbol{Q}$, which will be referred to by Φ_{rec}, differs from the azimuth of the true reaction plane, $\Phi_{r.p.}$, by

$$\Delta\Phi = \Phi_{r.p.} - \Phi_{rec}. \tag{3.8}$$

This is shown schematically in Figure 3.6b. It is important to know $\Delta\Phi$, i.e., the accuracy with which the azimuth of the reaction plane is determined. In particular, the value of $<\cos(\Delta\Phi)>$ is used for various corrections [68].

In this work, a method proposed in [69] is followed in order to estimate the value of $<\cos(\Delta\Phi)>$. Events are randomly divided into two subevents. For each of the subevents, the vector $\boldsymbol{Q}$ and the azimuth Φ_{rec} are determined. Figure 3.7a shows the distribution of

$$\Phi^{1-2} = \Phi^1_{rec} - \Phi^2_{rec} \tag{3.9}$$

for the MedB event sample in the Ru+Ru experiment. A typical gaussian shape on a constant offset can be observed. This is due to the fluctuations in the number of



Figure 3.6: *A schematic view of a semi-central collision as seen from a side (a) and in the transverse plane (b). In panel (b), the beam axis is orthogonal to the figure.* $\boldsymbol{b}$ *is the impact parameter vector.* $\boldsymbol{Q}$ *is reconstructed from* $\boldsymbol{p}_t$ *vectors of the reaction products following Eq. 3.7.* $\Delta\Phi$ *is the difference of the azimuthal angles of* $\boldsymbol{b}$ *and* $\boldsymbol{Q}$*. The picture in panel (b) is adapted from [69].*

particles and their $\boldsymbol{p}_t$ vectors, from which Φ^1_{rec} and Φ^2_{rec} are calculated. Figure 3.7b shows the dependence of the dispersion of Φ^{1-2} distribution on collision centrality given by TMUL. For the peripheral collisions (low TMUL), $RMS(\Phi^{1-2})$ decreases with increasing TMUL. This indicates that the determination of $\Phi_{r.p.}$ improves, which is due to the increasing number of particles used for the estimation of vectors $\boldsymbol{Q}_1$ and $\boldsymbol{Q}_2$. On the other hand, for very central reactions (high TMUL), $RMS(\Phi^{1-2})$ increases again because the reaction plane itself is not well defined.

It was proven that, if the fluctuations of the $\boldsymbol{Q}$ vector are gaussian (which is true when the number of particles used to calculate $\boldsymbol{Q}$ is large), the value of $<\cos(\Delta\Phi)>$ depends on $<\cos(\Phi^{1-2})>$ as shown by the line in Figure 3.7c. Therefore, for any considered event sample, the correction factor $<\cos(\Delta\Phi)>$ can be determined if the value of $<\cos(\Phi^{1-2})>$ is known. The values of $<\cos(\Phi^{1-2})>$ and the resulting values



Figure 3.7: *(a) The distribution of $\Phi^1_{rec} - \Phi^2_{rec}$ (see text for definition) for the medium-bias event sample in the Ru+Ru experiment. (b) The dependence of the dispersion RMS(Φ^{1-2}) on collision centrality given by TMUL. (c) The dependence of the correction factor $<cos(\Delta\Phi)>$ on $<cos(\Phi^{1-2})>$ [69].*

TMUL	50 – 85	85 – 95	95 – 140	all
$<\cos(\Phi^{1-2})>$	0.43	0.38	0.30	0.38
$<\cos(\Delta\Phi)>$	0.80	0.77	0.70	0.77

Table 3.2: *The values of $<cos(\Phi^{1-2})>$ and $<cos(\Delta\Phi)>$ for event categories of different centralities given by TMUL.*

of the correction factors $<\cos(\Delta\Phi)>$ calculated for three event categories considered in this work are listed in Table 3.2.

Chapter 4

Distributions of pions, protons and deuterons

In this chapter, the characteristics of the most abundant products of Ru+Ru collisions at 1.69 A GeV beam kinetic energy, i.e., pions, protons and deuterons will be discussed. The analysis concentrates on the particles that are registered by the CDC. The results will be used as a reference to the measurements on kaons presented in the following part of this work.

4.1 Identification

For positively charged particles, the identification is based on the mass parameters ($Mass_{CDC}$) assigned to particle tracks. In Figure 2.4b the distribution of $Mass_{CDC}$ is shown for particles with momenta lower than 3 GeV/c. The identification conditions are set according to the minima observed in the mass spectrum:

$$\begin{aligned} Mass_{CDC} &< 0.5\ \mathrm{GeV/c^2}, && \text{for } \pi^+\text{mesons}, \\ 0.5\ \mathrm{GeV/c^2} < Mass_{CDC} &< 1.5\ \mathrm{GeV/c^2}, && \text{for protons}, \\ 1.5\ \mathrm{GeV/c^2} < Mass_{CDC} &< 2.3\ \mathrm{GeV/c^2}, && \text{for deuterons}\ . \end{aligned} \tag{4.1}$$

With the above conditions particle species can be separated quite reliably. Nevertheless, the yields are not free from background contamination, which is estimated to be less than 10%, even for the highest particle momenta considered.

Since π^- mesons are practically the only negatively charged reaction products, the yield of π^- mesons should be free of any background contamination. All particles of which trajectories are bent in the opposite direction to proton trajectories are considered to be π^- mesons.